



PLANT GROWTH  
REGULATOR

# CutDown™

Plant Growth Regulator for Turfgrass

**ACTIVE INGREDIENT**

Paclobutrazol: ((±)-(R\*,R\*)-b-[(4-chlorophenyl) methyl]-a-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol):..... 22.3%

**OTHER INGREDIENTS:**..... 77.7%

**TOTAL**.....100.0%

CutDown™ contains 2 lbs. active ingredient per gallon.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

**FIRST AID**

<b>IF SWALLOWED</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF ON SKIN OR CLOTHING</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF INHALED</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>

**HOT LINE NUMBER**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact **1-888-875-1724 for emergency medical treatment information.**

# PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist.

## PERSONAL PROTECTIVE EQUIPMENT

### Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category F on an EPA chemical resistance category selection chart.

### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride, or viton
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Wash PPE after each day's use.

## USER SAFETY RECOMMENDATIONS

### Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately, if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

## PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

## DIRECTIONS FOR USE

**Failure to follow directions and precautions on this label may result in plant injury or poor disease control.**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Read all label directions carefully before use.

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride, or viton
- Shoes plus socks

## GENERAL INFORMATION

CutDown is a systemic plant growth regulator that slows vegetative growth by inhibiting gibberellin biosynthesis for use on turfgrass and trees. For turf, when CutDown is applied as directed, vertical growth of turf will be slowed within 3-10 days, resulting in reduced mowing frequency for a 6- to 8- week period. Following CutDown applications, turf will gradually undergo increased greening and density, which may persist up to 12 weeks.

## TURF

CutDown, a specialty plant growth regulator that slows vertical grass growth and reduces mowing for up to two (2) months after application. The frequency of mowing can be reduced by up to 50% during the period of effective retardation. Use of CutDown on fine turf should be accompanied by moderate-to-high fertility to maintain turfgrass appearance and reduce discoloration.

## USE DIRECTIONS

Turf	Application Rate (Product amount / a.i.- lbs per acre)	Application(s) May Result
St Augustine, Bermuda & similar warm season grasses	0.5 lbs. a.i. 32 fl. oz prod/A (Sandy soil)  0.75 lbs a.i. 48 fl. oz prod/A (Clay soil)	Reduced mowing for 6-8 weeks on established hybrid bermudagrass, bent grass and perennial ryegrass fairways, tees and roughs and on St. Augustinegrass and Kentucky bluegrass/perennial ryegrass turf.  Reduce potential for scalping and increase turf density and tighter-knit of all turfgrass areas.
Bluegrass, Rye, Fescue, Bent & similar cool season grasses (except putting greens)	0.25 to 0.50 lbs a.i. 16 -32 fl oz prod/A  Reduce by 50% if <i>Poa annua</i> is a major turfgrass population	Suppression of <i>Poa annua</i> by reducing its growth and its compatibility, leading to selective control after prolonged use.
Bentgrass Putting Greens & Overseeded Bermuda	0.10 to 0.25 lbs a.i. 6.4 -16 fl oz prod/A	

Apply CutDown with standard pressurized application equipment with by-pass or mechanical agitation using 50 mesh strainer to turf in sufficient amount of water (minimum 1 gal water /1,000 sq ft.= 43.5 gals/A) to ensure spray coverage uniformly to turf and to avoid skipping and/or overlapping.

For best results, use 2 to 5 gals water /1,000 sq ft. and apply 0.25 inch of water within 24 hours after application to remove product from foliage and onto soil surface.

CutDown can be used any time when established hybrid bermudagrass and St. Augustinegrass are green, actively growing and have recovered from dormancy after winter.

CutDown can be applied in spring after green-up and after turf has been mowed once or twice. Apply at least 1 month before onset of high air temperatures. In late summer/early fall, apply at least 1 month before anticipated first killing frost.

Apply anytime after overseeded turf has established itself successfully. Do not apply after March 15th to avoid delay in Bermudagrass green-up.

Maintain moderate soil moisture conditions before and after each application to achieve best plant growth retardation effect.

A repeat application of CutDown within the same growing season may be made 8 weeks apart for growth regulation & 4 to 6 weeks apart for color, quality enhancement after the initial application. Do not make more than 3 applications per calendar year. Do not apply more than 4 quarts per acre per year (2 lbs. a.i./A).

CutDown will not prevent seedhead production if used alone. A tank mixture for a spray program using a seed head control agent (Embark®) is required when seedhead visibility is a problem. If Embark is used for *Poa annua* seedhead control, apply CutDown at least 14 days after the Embark application.

### Weed, Insect and Diseases:

If crabgrass or other annual grassy weeds have been a problem in the past, an application of an appropriate preemergence weed control product should be made before the use of CutDown. Space applications of CutDown and preemergence product at least 4 weeks apart.

Broadleaf weeds are not significantly affected by CutDown. To control the growth of weeds, treat with an appropriate herbicide when weeds are actively growing. Carefully follow label directions.

If a weed, disease, or insect problem occurs after application of CutDown, apply control product at the recommended rate. CutDown is compatible with most existing control products. A compatibility test should be conducted with product(s) with which you do not have prior experience.

### Regulator response and Turf cultivar

Excessive irrigation and/or nitrogen fertilization may shorten period of growth regulation. Regulator response will vary somewhat according to turf variety. For St. Augustinegrass, Bitter Blue will be the most responsive and Floratam the least responsive.

### HYBRID BERMUDAGRASS

Sensitivity/Activity	Cultivar	Period of Growth Regulation	Period of Color Response
Medium/Good	Tifway I, II	5 - 6 weeks	6 - 8 weeks

### ST. AUGUSTINEGRASS

Sensitivity/Activity	Cultivar	Period of Growth Regulation	Period of Color Response
Low/Moderate	Floratam, Florawlawn	4 - 6 weeks	5 - 8 weeks
Medium/Good	Floratine, Raleigh, Texas, Common	6 - 7 weeks	7 - 8 weeks
High/Excellent	Bitter Blue, Seville	7 - 8 weeks	8 - 10 weeks

## GENERAL PRECAUTIONS

**Not** for use on bermudagrass putting greens except for winter overseeding enhancement use.

**Not** for use on athletic fields under heavy traffic where maximum growth potential of turf is desired.

**Not** for use on shrubs, flowers, fruits, or vegetable plants (applications to turf areas under trees will not affect/harm trees).

**Do not** use during periods of extreme dry or cold weather conditions, or during heavy insect or disease activity.

**Do not** apply product when soil is already saturated. Heavy rainfall or irrigation in the treated areas may cause active ingredient to move laterally on slopes and collect in low areas. These areas may undergo more severe growth control for a longer period of time.

Do not use on areas containing greater than 70% *Poa annua*, since discoloration of *Poa* may be unacceptable.

Delay treatment of newly-sodded or sprigged turf until grass has knitted down and rooted firmly.

Delay sprigging for at least 4 weeks and sodding at least 2 weeks after application is made.

Withhold application on large turf areas that have been thinned from winter drainage, disease, or insects until desired fill-in is achieved.

Assure that dosage rates are measured accurately since rates greater than those recommended may cause undesirable turf growth control and may discolor areas temporarily.

**Do not** mow treated grass for at least three days following application.

**Do not** use on areas to be cultivated for food or food crops or to be resown with grasses within two years of treatment.

**Do not** apply more than 4 qts. per acre per year (2 lbs. a.i./A).

**Do not** graze treated areas or harvest for forage or hay.

**Do not** apply this product through any type of irrigation equipment.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

**PESTICIDE STORAGE:** Keep container closed when not in use. Do not store near food or feed. Protect from freezing. In case of spill or leak on floor or paved surfaces, soak up with sand, earth or synthetic absorbent. Remove to chemical waste area.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the hazardous waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

## LIMITED WARRANTY AND DISCLAIMER

Phoenix Environmental Care LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below.

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such use or the results to be obtained if not used in accordance with printed directions and established safe practice. To the fullest extent permitted by law, buyer's exclusive remedy and manufacturer's or seller's exclusive liability for any and all claims, losses, damages or injuries resulting from the use or handling of this product, whether or not based in contract, negligence, strict liability in tort or otherwise shall be limited, at the manufacturer's option to replacement of, or the repayment of the purchase price for, the quantity of product with respect to which damages are claimed.

### Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Phoenix Environmental Care LLC or the seller. All such risks shall be assumed by buyer.

### Terms and Conditions of Use

If terms of the Limited Warranty & Disclaimer and Inherent Risks of Use are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer and Inherent Risks of Use.

CutDown™ is a trademark of Phoenix Environmental Care, LLC

Embark® is a trademark of PBI/Gordon Corp

(Made In China)

CPC 88674  
1020-01-7/12/10