

Nufarm

Propiconazole SPC 14.3 MEC

Fungicide

Nufarm Propiconazole SPC 14.3 MEC Fungicide contains propiconazole, the active ingredient used in Banner MAXX® and Alamo®.



Put the Power of Prevention to Work with Nufarm Propiconazole SPC 14.3 MEC

When it comes to disease control, preventive and systemic control are two of the most important attributes to look for in a fungicide, and two that Nufarm Propiconazole SPC 14.3 MEC Fungicide delivers. Propiconazole provides excellent preventive, systemic activity against many of the worst diseases in turf, including anthracnose, brown patch, dollar spot, take-all patch, necrotic ring spot, summer patch and *Fusarium*. In ornamentals and trees, Nufarm Propiconazole SPC 14.3 MEC controls powdery mildews, rusts, leafspots, scabs, and blights. Plus, Nufarm Propiconazole features a high-quality formulation for hassle-free application and outstanding tank-mix compatibility.

KEY PERFORMANCE BENEFITS

Broad-spectrum control across turf and ornamentals to provide excellent, dependable control

Ideal tank-mix compatibility and easy mixing into a clear solution

Offers systemic activity with long-lasting control

Low-odor, microemulsion concentrate liquid

Preventive control for up to 21 days



ACTIVE INGREDIENT

propiconazole

% ACTIVE INGREDIENT

14.30%

FORMULATION

microemulsion concentrate (MEC)

CHEMICAL FAMILY

triazole

GROUP NUMBER

3

EPA REGISTRATION

228-623

BRAND ALTERNATIVE

Banner MAXX® and Alamo®

MODE OF ACTION

sterol biosynthesis in membranes

RESTRICTED USE

no

SIGNAL WORD

warning

PACKAGING INFORMATION

4 x 1 gal
2 x 2.5 gal

LABELED USES

Turf
Ornamentals
Trees
(Flare root injection)*

DISEASES CONTROLLED

TURF:

Anthrachnose
Brown patch
Dollar spot
Fusarium
Gray leaf spot
Gray snow mold
Leafspot
Necrotic ring spot
Pink patch
Pink snow mold
Powdery mildew
Red thread
Rust
Spring dead spot
Stripe smut
Summer patch
Take-all patch
Yellow patch
Zoysia patch

ORNAMENTAL:

Conifer blights
Flower blight
Leaf blights/spots
Alternaria spp
Brown leaf spot
Greasy spot
Septoria spp
Anthrachnose
Downy spot
Scab
Powdery mildew
Rust

TREES:

Oak wilt
Dutch elm disease
Sycamore anthracnose
Leaf diseases of crab apple

SEE LABEL FOR COMPLETE LIST AND SPECIFICATIONS OF DISEASES CONTROLLED

Nufarm

Propiconazole SPC 14.3 MEC

Fungicide

APPLICATION RECOMMENDATIONS

TURF

Apply 0.5-4 fl oz/1000 sq ft or 22-176 fl oz/A

Use in a preventive disease control program. Apply after mowing OR allow sprayed area to completely dry before mowing. For control of soil-borne diseases, this product can be watered in after application. For control of foliar diseases, allow sprayed area to completely dry before irrigation. Refer to product label for specific rate information.

NURSERIES (FIELD) AND LANDSCAPE PLANTINGS

Use 2-24 fl oz/100 gallons of water for control of diseases of ornamental plant species. Refer to product label for specific rate information.

For outdoor uses, apply up to 5.4 gallons of this product/acre/crop/calendar year. For general disease control in landscapes, apply 6-8 fl oz/100 gallons of water every 21 days. For best control, begin applications before disease development.

TREES

Refer to the Nufarm Propiconazole SPC 14.3 MEC label for instructions on tree preparation and rates for specific diseases.

Better Choices. Better Business. | www.nufarm.com/us

For specific application rates, directions, mixing instructions and precautions, read the product label. Please visit www.nufarm.com/us to download a full label.

©2010 Nufarm. Important: Always read and follow label instructions.
Banner MAXX® and Alamo® are trademarks of a Syngenta Group Company.

10-TO-0039



*Not registered in California