



Quincept[®]

Herbicide

Quincept[®] contains 2,4-D, quinclorac and dicamba.



Control Crabgrass and 200 More Weeds with the Power of Quincept[®]

Quincept[®] herbicide from Nufarm offers the perfect balance of power and performance to control both grass and broadleaf weeds. Quincept combines the ideal ratio of active ingredients to deliver unbeatable results on crabgrass, foxtail, barnyardgrass, signalgrass and more than 200 broadleaf weeds.

Use Quincept in the spring as a clean-up product for areas that missed a pre-emerge grass herbicide application or late season for weed escapes. Quincept delivers the versatility you need to address many weed problems.

KEY PERFORMANCE BENEFITS

Post-emergent control of more than 200 tough broadleaf weeds plus crabgrass, foxtail, barnyardgrass and signalgrass

Three active ingredients pack a powerful punch

Perfect late season for weed escapes

Ideal for broadcast spray or spot applications

ACTIVE INGREDIENTS

2,4-D, quinclorac
and dicamba

% ACTIVE INGREDIENTS

13.24%, 8.25% and 1.38%

FORMULATION

liquid

CHEMICAL FAMILY

phenoxy, quinoline
carboxylic acid
and benzoic acid

GROUP NUMBER

4

EPA REGISTRATION

228-531

MODE OF ACTION

synthetic auxin
growth regulator

RESTRICTED USE

no

SIGNAL WORD

caution

PACKAGING INFORMATION

24 x 8 oz
8 x 64 oz
2 x 2.5 gal



LABELED USES

Residential and non-residential turfgrasses (*grounds or lawns around residential and commercial establishments, multi-family dwellings, military and other institutions, parks, airports, roadsides, schools, picnic grounds, athletic fields, houses of worship, cemeteries, golf courses and sod farms*)

WEEDS CONTROLLED

BROADLEAF WEEDS:	Dock	Oxeye daisy	Tall nettle
Alder	Dollar weed	Parsley-piert	Thistle
Annual yellow sweet clover	Dogbane	Parsnip	Tick trefoil
Artichoke	Dogfennel	Pearlwort	Toadflax
Austrian fieldcress	Elderberry	Pennycress	Trailing crownvetch
Bedstraw	English daisy	Pennywort	Tumble mustard
Beggartick	Fall dandelion	Peppergrass	Tumble pigweed
Biden	False dandelion	Pepperweed	Tumbleweed
Bindweed	False flax	Pigweed	Velvetleaf
Bird vetch	False sunflower	Pineywoods bedstraw	Venice mallow
Bitterweed	Fiddleneck	Plains coreopsis	Veronica
Bitter wintercress	Field bindweed	Plantain	Vervain
Black-eyed Susan	Field pansy	Poison hemlock	Vetch
Black medic	Fleabane (<i>daisy</i>)	Poison ivy	Violet, wild
Black mustard	Flixweed	Poison oak	Virginia buttonweed
Black-seed plantain	Florida betony	Pokeweed	Virginia creeper
Blessed thistle	Florida pusley	Poorjoe	Virginia pepperweed
Blue lettuce	Frenchweed	Povertyweed	Wavyleaf bullthistle
Blue vervain	Galinsoga	Prostrate spurge	Western clematis
Boxelder	Garlic mustard	Prostrate vervain	Western salsify
Bracted plantain	Goathead	Puncture vine	White clover
Brassbuttons	Goatsbeard	Purslane	White mustard
Bristly oxtongue	Goldenrod	Ragweed	Wild aster
Broadleaf dock	Ground ivy	Red clover	Wild buckwheat
Broadleaf plantain	Gumweed	Redroot pigweed	Wild carrot
Broomweed	Hairy bittercress	Red sorrel	Wild four-o'clock
Buckhorn	Hairy fleabane	Redstem filaree	Wild garlic
Buckhorn plantain	Hawkweed	Rough cinquefoil	Wild geranium
Bulbous buttercup	Healall	Rough fleabane	Wild lettuce
Bull nettle	Heartleaf drymary	Roundleafed marigold	Wild marigold
Bull thistle	Hedge bindweed	Rush	Wild onion
Burdock	Hedge mustard	Russian pigweed	Wild parsnip
Burning nettle	Hemp	Russian thistle	Wild radish
Bur ragweed	Henbit	St. Johnswort	Wild rape
Burweed	Hoary cress	Scarlet pimpernel	Wild strawberry
Buttercup	Horsetail	Scotch thistle	Wild sweet potato
Canada thistle	Indiana mallow	Sheep sorrel	Wild vetch
Carolina geranium	Ironweed	Shepherd's-purse	Willow
Carpetweed	Jewelweed	Slender plantain	Witchweed
Catchweed bedstraw	Jimsonweed	Smallflower galinsoga	Woodsorrel
Catsear	Kochia	Smartweed	Woolly croton
Catnip	Knawel	Smooth dock	Woolly morningglory
Chickweed	Knotweed	Smooth pigweed	Woolly plantain
Chicory	Lambsquarters	Sneezeweed	Wormseed
Cinquefoil	Lespedeza	Southern wild rose	Yarrow
Common chickweed	Locoweed	Sowthistle	Yellow rocket
Common mullein	Lupine	Spanishneedle	Yellowflower pepperweed
Common sowthistle	Mallow	Spatterdock	and other broadleaf weeds
Corn chamomile	Marshelder	Speedwell	
Creeping Jenny	Matchweed	Spiny amaranth	
Crimson clover	Mexicanweed	Spiny cocklebur	
Croton	Milk vetch	Spotted catsear	
Cudweed	Milkweed	Spotted knapweed	
Curly dock	bloodflower	Spotted spurge	
Curly indigo	Mugwort	Spurge	
Dandelion	Morningglory	Spurweed	
Dead nettle	Mousear chickweed	Stinging nettle	
	Musk thistle	Stinkweed	
	Mustard	Stitchwort	
	Narrowleaf plantain	Strawberry clover	
	Narrowleaf vetch	Sumac	
	Nettle	Sunflower	
	Orange hawkweed	Sweet clover	
	Oxalis		



Herbicide

APPLICATION RECOMMENDATIONS

For single application, use 7 to 8 pts per acre or 112 to 128 ounces per acre. For spot treatments, apply 2.6 to 2.9 fluid ounces per 1,000 sq ft.

Ensure that spray volumes are adequate to completely cover weeds, especially when dense weed infestations make it difficult to completely cover the foliage of target weeds.

Early Quincept applications will not control weeds germinating later in the season so a second application may be necessary.

DUAL APPLICATIONS

A single application should control less mature, light weed populations; however, a second application may be required to control heavy, mature broadleaf weed infestations. Under certain conditions, annual grasses in the 2 to 4-tiller stages may require a sequential application. If a second application is required, allow 14 to 21 days between applications.

Better Choices. Better Business.
www.nufarm.com/us

For specific application rates, directions, mixing instructions and precautions, read the product label. Please visit www.nufarm.com/us to download a full label.

©2010 Nufarm. Important: Always read and follow label instructions. Quincept® is a trademark of Nufarm.
09-TO-0077

