

Nufarm

Quinclorac SPC 75 DF Herbicide

Nufarm Quinclorac SPC 75 DF Herbicide contains quinclorac, the active ingredient used in Drive®.



Tackle Tough Turf Weeds with Nufarm Quinclorac SPC 75 DF Herbicide

Trust Nufarm Quinclorac SPC 75 DF Herbicide for post-emergent weed control. Just a single application controls a broad-spectrum of grass and broadleaf weeds—including clover, crabgrass and foxtail—and keeps them down for up to 30 to 45 days. Plus, Quinclorac SPC can be used in a variety of turf species.

KEY PERFORMANCE BENEFITS

Controls hard-to-control weeds such as crabgrass, clover, speedwell, dandelion, foxtail and dollarweed

Delivers up to 30 to 45 days of residual control

Offers excellent seeding and overseeding flexibility—annual bluegrass, annual ryegrass, buffalograss, common Bermuda, hybrid Bermuda, tall fescue and zoysia any time before or after application

ACTIVE INGREDIENT

quinclorac

% ACTIVE INGREDIENT

75%

FORMULATION

dry flowable

CHEMICAL FAMILY

quinoline carboxylic acid

EPA REGISTRATION

228-592

MODE OF ACTION

synthetic auxin

GROUP NUMBER

4

RESTRICTED USE

no

SIGNAL WORD

caution

PACKAGING INFORMATION

6 x 1 lb case



LABELED USES

Residential and ornamental
turfgrasses

WEEDS CONTROLLED

GRASSY WEEDS:

Barnyardgrass
Crabgrass, large^{1,4}
Crabgrass, smooth^{1,4}
Foxtail, giant¹
Foxtail, green¹
Foxtail, yellow¹
Kikuyugrass^{2,3}
Signalgrass, broadleaf¹
Torpedograss³

BROADLEAF WEEDS:

Bindweed, field
Clover, hop
Clover, red
Clover, white
Dandelion, common²
Daisy, English^{2,5}
Dollarweed
Geranium, Carolina
Medic, black
Morningglory spp.
Speedwell, common
Speedwell, slender
Speedwell, thymeleaf
Violet, wild

¹ Ensure that spray volumes are adequate to completely cover weeds, especially when dense weed infestations make it difficult to completely cover the foliage of target weeds. Optimum control is achieved when applications of this product plus methylated seed oil is applied either before second tiller or as weed grasses mature.

² Tank mix partner or sequential application required.

³ Make 3 sequential applications of 0.245 ounce of product per 1,000 square feet at 14 to 21 day intervals.

⁴ Biotypes of large and smooth crabgrass in California have shown varied response to this product. If control failure occurs following the application of a full or split application rates, do not reapply this product and change to a herbicide with a different mode of action.

⁵ NOT FOR USE to control this weed in California.

Nufarm

Quinclorac SPC 75 DF

Herbicide

APPLICATION RECOMMENDATIONS

Broadcast: Apply 1 pound/A or 0.367 ounce/1,000 sq ft with 1.5 pints/A or 0.55 ounce/1,000 sq ft of methylated seed oil.

Spot treatment: 0.367 ounces per 1,000 sq ft of treated area with 1.5 pints/A or 0.55 ounce/1,000 sq ft of methylated seed oil.

Apply when weeds are actively growing and in early stages of growth. More mature grasses will be more difficult to control and may require a second application. Mature, drought-stressed grassy weeds will be more difficult to control so adequate soil moisture is preferred. Adverse or extreme environmental conditions such as poor soil conditions, high temperatures, drought and cultural conditions may affect the performance of this product.

Better Choices. Better Business. | www.nufarm.com/us

For specific application rates, directions, mixing instructions and precautions, read the product label. Please visit www.nufarm.com/us to download a full label.

©2010 Nufarm. Important: Always read and follow label instructions.
Drive® is a trademark of BASF Corporation.

10-TO-0002

